

# Alcohol, DNA Methylation, and Cancer

**Marta Varela-Rey, Ph.D.; Ashwin Woodhoo, Ph.D.; Maria-Luz Martinez-Chantar, Ph.D.; José M. Mato, Ph.D.; and Shelly C. Lu, M.D.**

**Marta Varela-Rey, Ph.D.**, is a CIBERehd postdoctoral fellow, **Ashwin Woodhoo, Ph.D.**, is a Ramón y Cajal postdoctoral fellow, and **Maria-Luz Martinez-Chantar, Ph.D.**, is group leader, all at the Centro de Investigación Cooperativa en Biosciencias (CIC bioGUNE), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBERehd), Technology Park of Bizkaia, Derio, Spain.

**José M. Mato, Ph.D.**, is director of CIC bioGUNE and of the Center for Cooperative Research in Biomaterials (CIC bioMAGUNE), as well as research professor and group leader at CIC bioGUNE, (CIBERehd), Technology Park of Bizkaia, Derio, Spain.

**Shelly C. Lu, M.D.**, is professor of medicine, associate chief of the Division of Gastrointestinal and Liver Diseases, and associate director of the USC Research Center for Liver Diseases, Keck School of Medicine, University of Southern California, Los Angeles, California.

Cancer is one of the most significant diseases associated with chronic alcohol consumption, and chronic drinking is a strong risk factor for cancer, particularly of the upper aerodigestive tract, liver, colorectum, and breast. Several factors contribute to alcohol-induced cancer development (i.e., carcinogenesis), including the actions of acetaldehyde, the first and primary metabolite of ethanol, and oxidative stress. However, increasing evidence suggests that aberrant patterns of DNA methylation, an important epigenetic mechanism of transcriptional control, also could be part of the pathogenetic mechanisms that lead to alcohol-induced cancer development. The effects of alcohol on global and local DNA methylation patterns likely are mediated by its ability to interfere with the availability of the principal biological methyl donor, S-adenosylmethionine (SAME), as well as pathways related to it. Several mechanisms may mediate the effects of alcohol on DNA methylation, including reduced folate levels and inhibition of key enzymes in one-carbon metabolism that ultimately lead to lower SAME levels, as well as inhibition of activity and expression of enzymes involved in DNA methylation (i.e., DNA methyltransferases). Finally, variations (i.e., polymorphisms) of several genes involved in one-carbon metabolism also modulate the risk of alcohol-associated carcinogenesis. **KEY WORDS: Alcohol consumption; chronic alcohol use; alcohol use disorders; alcohol-induced cancer development; risk factors; epigenetics; epigenetic mechanisms; cancer; upper aerodigestive tract cancer; colorectum cancer; breast cancer; carcinogenesis; DNA methylation**

**A**ccording to the World Health Organization (WHO) Global Burden of Disease Project, alcohol accounts for approximately 3.2 percent of all deaths per year worldwide (corresponding to 1.8 million people) and is causally related to more than 60 different medical conditions (Rehm et al. 2004). Cancer formation (i.e., carcinogenesis) is one of the most significant consequences attributed to alcohol consumption, and approximately 3.6 percent of all cancer-related cases (5.2 percent in men and 1.7 percent in women) worldwide, as well as 3.5 percent of all cancer-related deaths are related to chronic alcohol drinking (Boffetta et al. 2006). Based on available epidemiological data, an international group of epidemiologists and alcohol researchers concluded that alcohol induces carcinogenesis in numerous organs,

including the upper aerodigestive tract, liver, colorectum, and female breast (Baan et al. 2007).

Several pathogenic mechanisms contribute to alcohol-induced carcinogenesis in each type of cancer. The most commonly cited mechanisms include the effect of acetaldehyde—the first metabolite of ethanol oxidation—and oxidative stress (Seitz and Stickel 2007). Increasing evidence shows that alcohol also can induce epigenetic alterations, for example, in pathological conditions such as fetal alcohol spectrum disorders (Kobor and Weinberg 2011). Epigenetic alterations also are a hallmark of cancer development in general (Esteller 2008). Therefore, this article reviews the available evidence that such changes also could be an important contributory factor to alcohol-induced carcinogenesis. In particular, it discusses the role of DNA

methylation in carcinogenesis and how alcohol may affect the pathways that regulate the availability of S-adenosylmethionine (S-AdoMet), the principal biological methyl donor for methylation reactions. (For a list of abbreviations of the names for genes, proteins, and other compounds as well as their functions, see table).

## DNA Methylation and Cancer

DNA methylation plays a critical role in the control of gene activity. This methylation almost exclusively involves the addition of a methyl group to carbon 5 of cytosine nucleotides, and specifically those cytosines that precede guanines (i.e., are part of CpG dinucleotides). CpG dinucleotides tend to cluster either in regions called CpG islands, which are located in approximately 60 percent of human gene promoters, or in regions that contain large repetitive DNA sequences (e.g., centromeres and retrotransposons). In the former case, the CpG dinucleotides generally tend to remain unmethylated, whereas in the latter case they mostly are methylated to prevent chromosome instability (Rodriguez-Paredes and Esteller 2011) (figure 1). DNA methylation also occurs in CpG island shores—that is, regions of lower CpG density that are located close to CpG islands (i.e., within 2 kb). In these three cases, DNA methylation typically is associated with repression of gene expression (i.e., transcription). In a few cases, however, DNA methylation also can enable gene transcription, namely when the methylation occurs in the body (i.e., coding sequences) of the gene rather than the promoter (Portela and Esteller 2010).

DNA methylation is mediated (i.e., catalyzed) by three main enzymes called DNA methyltransferases (DNMTs), all of which transfer a methyl group from S-AdoMet to DNA:

- DNMT1, which also is referred to as the maintenance DNMT, completes the methylation of the par-

tially methylated (i.e., hemimethylated) DNA that is present in the cell after cell division and the accompanying DNA replication.

- DNMT 3A and DNMT 3B are referred to as de novo methyltrans-

ferases because they target unmethylated CpGs.

Aberrant epigenetic regulation, including altered DNA methylation, characterizes a wide range of diseases, including cancer (Portela and Esteller

**Table 1** Abbreviations of Gene Names, Protein Names, and Other Molecules


Abbreviation	Spelled-out Name
<b>Genes</b>	
<i>APC-1A</i>	Adenomatous polyposis coli
<i>BRCA1</i>	Breast cancer 1, early onset
<i>CHRNA3</i>	Cholinergic receptor, nicotinic, alpha 3
<i>CDKN2A</i>	Cyclin-dependent kinase inhibitor 2A
<i>p14ARF</i>	Cyclin-dependent kinase inhibitor 2A, isoform 4
<i>CDKN2B, p15INK4b</i>	Cyclin-dependent kinase 4 inhibitor B
<i>CDKN2A, p16Ink4A</i>	Cyclin-dependent kinase inhibitor 2A
<i>ER-α</i>	Estrogen receptor
<i>GATA2</i>	GATA binding protein 2
<i>GSTP1</i>	Glutathione S-transferase pi 1
<i>HOX2A</i>	homeobox B5
<i>MGMT</i>	O-6-Methylguanine-DNA methyltransferase
<i>hMLH1</i>	MutL homolog 1
<i>RASSF1A</i>	Ras association (RalGDS/AF-6) domain family member 1
<i>RB</i>	Retinoblastoma
<b>Proteins</b>	
BHMT	Betaine homocysteine methyltransferase
DHFR	Dihydrofolate reductase
DNMTs	DNA methyltransferases
LINE family member L1, LINE-1	Long interspersed nucleotide element-1
MAT	Methionine adenosyltransferase
MTR	Methionine synthase
MTHFR	methylenetetrahydrofolate reductase
MT	Methyltransferase
SFRP1	Secreted frizzled-related protein 1
TS	Thymidylate synthase
<b>Other Compounds</b>	
DHF	Dihydrofolate
dTMP	Deoxythymidine monophosphate
dUMP	Deoxyuridine monophosphate
Hcy	Homocysteine
Met	Methionine
5,10-MTHF	5,10-Methylenetetrahydrofolate
5-MTHF	5-Methyltetrahydrofolate
SAH	S-adenosylhomocysteine
S-AdoMet	S-adenosylmethionine
THF	Tetrahydrofolate

2010). Compared with normal cells, the epigenome of cancer cells shows profound changes in DNA methylation patterns as well as histone modifications patterns (Rodriguez-Paredes and Esteller 2011), including the following:

- Genome-wide hypomethylation (about 20 to 60 percent less overall 5-methyl-cytosine compared with normal cells); global loss of DNA methylation occurs at many genomic regions, such as repetitive elements and retrotransposons, resulting in chromosomal instability, and activation of transposable elements and endoparasitic sequences (e.g., LINE family member L1) (figure 1).
- Hypomethylation at specific promoters, resulting in aberrant activation of cancer-inducing genes (i.e., oncogenes) (e.g., *SERPINB5*) and inducing loss of imprinting at other DNA sites (e.g., *IGF2*).
- Hypermethylation of CpG island promoters of genes involved in central cellular pathways, such as DNA repair (e.g., at genes called *hMLH1*, *MGMT*, and *BRCA1*) and cell cycle control (e.g., at genes called *p16INK4a*, *p15INK4b*, and *RB*), resulting in the silencing by is the most recognized epigenetic disruption in human tumors.
- Aberrant DNA methylation at CpG island shores (e.g., in genes called *HOX2A* and *GATA2*) (Portela and Esteller 2010; Rodriguez-Paredes and Esteller 2011).

## One-Carbon Metabolism

Persuasive evidence indicates that several dietary factors, such as alcohol, can modulate DNA methylation patterns and increase susceptibility to disease, including cancer, by altering one-carbon metabolism (Choi and Friso 2010). The term one-carbon metabolism refers to a network of biochemical reactions in which a chemical unit containing one carbon atom (e.g., a methyl group) is transferred through several steps from a donor to another compound, such as DNA (figure 2). The first step in this chain of events is the transfer of the one-carbon group to tetrahydrofolate (THF), resulting in the formation of 5,10-MTHF. This molecule then can be used in the synthesis of the nucleotide thymidine in a



**Figure 1** DNA methylation patterns in normal and cancer cells. (a) Repetitive sequences generally are methylated at cytosine nucleotides in normal cells. Global loss of methylation in cancer cells leads to chromosomal instability and activation of endoparasitic sequences. (b) CpG islands in promoter sequences typically are unmethylated in normal cells whereas they can become hypermethylated in cancer cells, leading to transcriptional repression. Examples of genes affected are shown on the right. (c) Similar patterns are seen in CpG island shores, located in front (i.e., upstream) of promoters. (d) CpGs located in gene bodies frequently are methylated in normal cells; this pattern is reversed in cancer cells, leading to initiation of transcription at several incorrect sites.

reaction catalyzed by the enzyme thymidylate synthase (TS). In this reaction, one carbon group of the 5,10-MTHF is transferred to a molecule called deoxyuridine monophosphate (dUMP), resulting in the formation of deoxythymidine monophosphate (dTMP) and dihydrofolate (DHF). This reaction is considered to be a limiting step in DNA synthesis and, more importantly, reduces dUMP levels, which can lead to breaks in both strands of the DNA molecules. Finally, the enzyme dihydrofolate reductase (DHFR) catalyzes the reduction of DHF back into THF (Mato et al. 2008; Van der Put et al. 2001).

In an alternative set of reactions, 5,10-MTHF also can be converted to 5-methyltetrahydrofolate (5-MTHF) in a reaction catalyzed by the enzyme


methylenetetrahydrofolate reductase (MTHFR). The 5-MTHF generated then can be used in the remethylation of homocysteine (Hcy) to methionine, which is catalyzed by methionine synthase (MTR). In the liver and kidney, Hcy also can be remethylated into methionine through a folate-independent reaction, catalyzed by betaine homocysteine methyltransferase (BHMT). BHMT requires betaine, a break-down product (i.e., metabolite) of choline for its activity.

Methionine, in turn, is an essential amino acid that, together with ATP, participates in the formation of SAdoMet, in a reaction catalyzed by methionine adenosyltransferase (MAT). SAdoMet is the principal methyl donor for numerous reactions, including protein, RNA, DNA, and histone methylation. The

formation of SAdoMet is a limiting step in DNA methylation. After transfer of its activated methyl group in the methylation reactions, SAdoMet is converted into S-adenosylhomocysteine (SAH). This compound also can affect DNA methylation because it is a potent competitive inhibitor of methyltransferases, including DNMTs. Both an increase in SAH levels and a decrease in the SAdoMet-to-SA-H ratio can inhibit transmethylation reactions (Lu 2000).

### Polymorphisms of One-Carbon Metabolism Enzymes

Several studies have shown that variations (i.e., polymorphisms) in several of the genes encoding enzymes involved in one-carbon metabolism and, consequently, in the resulting



**Figure 2** One-carbon metabolism with a schematic representation of the role of methionine in folate metabolism and transmethylation reactions and steps that are inhibited by alcohol. BHMT: betaine homocysteine methyltransferase; DHF: dihydrofolate; DHFR: dihydrofolate reductase; DNMT: DNA methyltransferase; dTMP: deoxythymidine monophosphate; dUMP: deoxyuridine monophosphate; Hcy: homocysteine; MAT: methionine adenosyl transferase; Met: methionine; MT: methyltransferase; 5-MTHF: 5-methyltetrahydrofolate; 5,10-MTHF: 5,10-methylenetetrahydrofolate; MTHFR: methylenetetrahydrofolate reductase; MTR: methionine synthase; SAH: S-adenosylhomocysteine; SAdoMet: S-adenosylmethionine; THF: tetrahydrofolate; TS: thymidylate synthase.

enzymes can impact the levels of various metabolites generated during these reactions. For some of these metabolites, altered levels in the body can be associated with either increased or decreased cancer susceptibility.

### **Polymorphisms in the Gene Encoding MTHFR.**

As mentioned above, MTHFR plays a central role in methionine formation by mediating the synthesis of 5-MTHF, which serves as a substrate for the remethylation of Hcy to methionine. Several polymorphisms in the *MTHFR* gene have been reported; two of these, known as the *C677T*<sup>1</sup> and *A1298C* *MTHFR* variants, in particular have been investigated in relation to cancer susceptibility. The *C677T* variant occurs in 8 to 15 percent of the population and results in a reduction of *MTHFR* activity. Thus, compared with people carrying two copies of the normal *MTHFR* gene (i.e., homozygous for *MTHFR C* [*CC*]), people carrying one normal gene copy and one variant gene copy (i.e., heterozygous with one *MTHFR C* and one *MTHFR T* [*CT*]) show a 45 percent reduction in *MTHFR* activity, and those homozygous for the *MTHFR T* variant (*TT*) show a 70 percent decrease of *MTHFR* activity (Motulsky 1996; Van der Put et al. 1995). Persons with the *TT* genotype have low levels of folate and vitamin B12 in the blood, as well as increased Hcy levels. Studies of the plasma folate and Hcy levels in people carrying one or two copies of the *A1298C MTHFR* polymorphism, however, have yielded inconsistent results (Sharp and Little 2004).

### **Polymorphisms in the Gene Encoding TS.**

*TS* catalyzes the conversion of dUMP into dTMP using 5,10-MTHF. Because dUMP can be misincorporated into new DNA strands during DNA replication,

which can cause breaks in the double-stranded DNA molecules, the actions of *TS* can reduce the occurrence of DNA double-strand breaks. Two polymorphisms in the *TS* gene may be associated with an increased cancer risk:

- The presence of a repeated sequence of 28 nucleotides (i.e., a 28-bp tandem repeat) at the beginning of the *TS* gene (i.e., in the *TS* 5'-untranslated enhanced region); this variant is referred to as *TSER*;
- A 6-bp deletion/insertion at the end of the *TS* gene (i.e., in the *TS* 3'-untranslated region); this variant is known as *TS1494del6*.

### **Polymorphisms in the Gene Encoding MTR.**

*MTR* catalyzes the remethylation of Hcy to methionine; its activity is essential to maintain adequate folate pools. Most studies suggest that plasma Hcy levels are lower in carriers of a variant of the *MTR* gene called *A2756G*. To function properly, *MTR* must be maintained in an active form; this is achieved by another enzyme called *MTR* reductase (*MTRR*), for which polymorphisms also exist. For example, people homozygous for the *A66G MTRR* polymorphism had elevated levels of Hcy, indicating that the resulting *MTRR* enzyme led to higher-than-normal *MTR* activity (Van der Put et al. 2001).

## **Alcohol and DNA Methylation**

As mentioned above, several pathogenetic mechanisms for alcohol-induced carcinogenesis have been described, including the effects of acetaldehyde and oxidative stress (Seitz and Stickel 2007). More recently, increasing evidence indicates that alcohol may induce epigenetic alterations, in particular aberrant DNA methylation patterns, which also could be important contributory factors to alcohol-induced carcinogenesis. For example, excessive alcohol use is associated with increased risk of

colon cancer, which is characterized by global DNA hypomethylation as well as hypermethylation of certain genes (see below). Several mechanisms have been described to date that could contribute to these DNA methylation changes in cancer. These generally are associated with modulation of the pathways that regulate the availability of SAME.

### **Alcohol and Lipotropes**

The term lipotropes denotes compounds that help catalyze the breakdown of fat molecules in the body. Lipotropic nutrients (e.g., methionine, choline, folate, and betaine) are important dietary methyl donors and cofactors that play key roles in one-carbon metabolism. Dietary lipotropes influence the availability of SAME and, consequently, may influence genomic DNA methylation patterns and the expression of multiple cancer-related genes. For example, methyl-deficient diets can induce the development of liver tumors (i.e., hepatocarcinogenesis) in rats by causing global and gene-specific hypomethylation (for a review, see Ross and Poirier 2002). Chronic alcoholics frequently suffer from malnutrition that results in depletion of lipotropes (Seitz and Stickel 2007). The lack of these nutrients in heavy drinkers could possibly result in an altered SAME production, leading to changes in DNA methylation.

Similarly, malnutrition in alcoholics leads to a severe deficiency of other cofactors of one-carbon metabolism, such as folate (see below), vitamin B6, and vitamin B12. Long-term dietary intake of vitamin B6 is inversely correlated with the risk of developing colorectal cancer in women—that is, inadequate vitamin B6 intake increases the risk of this cancer. This effect is aggravated by chronic consumption of alcohol (Larsson et al. 2005).

### **Alcohol and Folate Status**

Epidemiological studies have suggested that reduced folate levels in the body

<sup>1</sup>The name of the variant indicates that the amino acid cysteine (abbreviated as C), which normally is found at position 677 in the *MTHFR* molecule, is replaced by a threonine (abbreviated as T). The names of other gene variants can be interpreted similarly.

increase the risk of several types of cancer, including those of the upper aerodigestive tract, colon/rectum, and breast (Kim 2005). For example, heavy drinkers with low methionine and folate levels have a significantly increased relative risk (RR) for colorectal cancer compared with occasional drinkers with normal methionine and folate intake (Giovannucci et al. 1995). In chronic alcoholics, serum folate levels are significantly reduced compared with healthy subjects (Cravo et al. 1996), likely because folate absorption is reduced in these patients (Halsted et al. 1971). Polymorphisms of the *MTHFR* genes also lead to reduced folate levels, but their contribution to carcinogenesis is tissue-dependent and often contradictory. This is discussed in more detail below.

Two main mechanisms have been described that may explain the cancer-promoting effects of limited folate levels: increased DNA instability and aberrant DNA methylation patterns (Hamid 2012). Folate deficiency alters the balance of the pool of nucleotides needed for the synthesis of new DNA molecules, leading to dUMP accumulation. As a result, dUMP is misincorporated into new DNA molecules; this and the subsequent repair processes can lead to double-strand breaks in the DNA and chromosomal damage, ultimately resulting in cancer. The aberrant DNA methylation patterns associated with folate deficiency are the result of folate's role in one-carbon metabolism (see figure 2). As mentioned above, 5-MTHF can be used in the remethylation of Hcy to methionine, which in turn generates SAME. Many studies have shown that folate deficiency reduces SAME levels and the SAME-to-SAH ratio as well as increases SAH concentrations (Kim 2005), all of which might contribute to carcinogenesis.

### **Alcohol and MTR Activity**

Alcohol also can affect SAME, SAH, and Hcy levels by reducing MTR activity, which in turn results in decreased SAME levels and enhanced generation

of Hcy and SAH (Barak et al. 1996) (see figure 2). To compensate for this decrease in MTR activity, the activity of BHMT is induced after alcohol ingestion. However, after extended periods of alcohol exposure this alternate pathway cannot be maintained. This results in a decrease in the hepatocyte level of SAME, increases in SAH and Hcy levels, and a reduced SAME-to-SAH ratio. These effects may contribute to the reduced hepatic SAME levels observed in patients hospitalized for alcoholic hepatitis (Lee et al. 2004).

**The major risk factors for developing hepatocellular carcinoma are viral infection, chronic alcoholism, and exposure to toxic substances called aflatoxins.**

### **Alcohol and MAT Activity**

As mentioned earlier, alcohol also contributes to the generation of oxidative stress through various mechanisms (for a review, see Seitz and Stickel 2007). Increased oxidative stress, in turn, can, at least in the liver, inactivate the MAT I/III enzymes that convert methionine to SAME (see figure 2). This inactivation results from the covalent modification of a critical cysteine residue at position 121 by nitric oxide and hydroxyl radicals. MAT inhibition causes decreased SAME levels, leading to reduced methylation reactions. The relevance of this pathway was demonstrated by findings that patients with alcoholic cirrhosis exhibit decreased hepatic MAT activity and SAME formation (Lu and Mato 2008). In addition, patients with alcoholic hepatitis also show reduced expression of the *MAT1A* gene that encodes the MAT isoenzymes in normal liver, thereby contributing to lower hepatic SAME levels (Lee et al. 2004).

### **Alcohol and DNMT Activity**

Finally, both ethanol and its first breakdown product (i.e., acetaldehyde) can impact methylation patterns by altering DNMT activity. Thus, studies found that acetaldehyde can inhibit DNMT activity in vitro (Garro et al. 1991) and that alcohol reduced *DNMT* mRNA levels in rats treated with alcohol for 9 weeks (Bielawski et al. 2002). Similarly, studies in humans found that *DNMT3a* and *DNMT3b* mRNA levels were significantly reduced in patients with chronic alcoholism compared with healthy control subjects (Bonsch et al. 2006).

### **Alcohol, DNA Methylation, and Cancer**

As described above, alcohol can interfere with one-carbon metabolism in several ways, thereby potentially generating aberrant DNA methylation patterns. The following sections review the currently available evidence indicating that alcohol-mediated changes in DNA methylation profiles contribute to the four main alcohol-associated cancers.

#### **Liver Cancer**

Liver cancer (i.e., hepatocellular carcinoma [HCC]) is a major cause of cancer-related death worldwide. The major risk factors for developing HCC are viral infection (i.e., with the hepatitis B or C viruses), chronic alcoholism, and exposure to toxic substances called aflatoxins. Alcohol remains the major cause of liver-related disease and deaths in the United States (Moghe et al. 2011). Alcohol-induced aberrant DNA methylation has been well characterized as a pathogenetic mechanism contributing to liver disease. Significant global DNA hypomethylation is associated with HCC with several etiologies, including chronic alcoholism, which may result in malignant transformation through mechanisms such as loss of imprinting and chromosomal instability, as described above (Calvisi

et al. 2007; Hernandez-Vargas et al. 2010). Similar to other types of cancer, global DNA hypomethylation in HCC is accompanied by greater-than-normal methylation levels (i.e., hypermethylation) at certain CpG sites. For example, a study analyzing 1,505 CpG sites in the promoter regions of 807 cancer-related genes in HCC tissues with different underlying causes, including chronic alcohol consumption, demonstrated an altered methylation pattern in 94 genes. Furthermore, for specific subsets of genes significant associations existed between methylation patterns and tumor progression (i.e., stage of the tumor and grade of differentiation) and background (i.e., cirrhotic versus non-cirrhotic surrounding tissue) (Hernandez-Vargas et al. 2010). Genes exhibiting hypermethylation included *RASSF1*, *APC*, and *CDKN2A*, all of which have important functions in the liver (Tischoff and Tannapfel 2008). A more recent study (Lampert et al. 2011) also detected aberrant hypermethylation of several genes, including *RASSF1*, *GSTP1*, *MGMT*, and *CHRNA3*, in alcohol-associated HCC. Finally, mRNA expression of RB1, an important cell cycle regulator, is decreased as a result of promoter methylation (Edamoto et al. 2003).

The link between alcohol and the mechanisms leading to aberrant methylation has been well elucidated in HCC. Chronic alcoholic patients have reduced hepatic MAT activity resulting from both decreased expression of the *MAT1A* gene and inactivation of the MAT I/III proteins. Reduced MAT activity, in turn, leads to decreased SAME biosynthesis (Tsukamoto and Lu 2001), which may contribute to the severe loss of global DNA methylation in HCC. Similarly, micropigs fed ethanol show reduced hepatic MTR activity and SAME/SAH ratio (Halsted et al. 1996) and rats fed ethanol exhibit decreased SAME levels and global DNA hypomethylation.

Patients heterozygous or homozygous for the previously mentioned C677T *MTHFR* polymorphism have

been shown to have a lower risk of developing alcohol-related HCC, but not of HCC with other etiologies (Saffroy et al. 2004). Similarly, people carrying variant alleles of both *MTHFR* and *TS* genes had a statistically significant reduced risk of developing HCC (Yuan et al. 2007). However, these findings sharply contradict another study showing that male patients with alcoholic cirrhosis who were homozygous for the C677T *MTHFR* polymorphism had an increased risk of developing HCC (Fabris et al. 2009). How can these conflicting reports be explained? In principle, the C677T *MTHFR* polymorphism has both protective and cancer-promoting effects. On the one hand, it can have a protective effect because it leads to an increase in the supply of 5,10-MTHF, which can serve as a substrate in the conversion of dUMP to dTMP. This would reduce misincorporation of dUMP into DNA, preventing DNA double-strand breaks and chromosomal instability. On the other hand, the reduced activity of the MTHFR enzyme encoded by the C677T *MTHFR* polymorphism would lead to lower levels of 5-MTHF that is used for the remethylation of Hcy to methionine, which in turn generates SAME. Thus, the polymorphism would result in lower SAME levels and SAME/SAH ratio as well as increased SAH concentrations, which would then contribute to carcinogenesis.

The overall effects of the C677T *MTHFR* polymorphism can be determined when folate levels are taken into account. In situations of adequate folate supply, the levels of 5,10-MTHF increase, leading to a protective effect as described above, possibly counteracting the effects of reduced SAME. In folate-deficiency situations, however, the levels of both 5,10-MTHF and SAME would be reduced, leading to chromosomal instability that would be exacerbated by DNA hypomethylation induced by reduced SAME. This model is consistent with a study in colon cancer showing that patients with the CC genotype have significantly reduced

risk of cancer development in situations of adequate folate supply, but that this protection is absent in folate deficiency (Ma et al. 1997). It also can reconcile the contradictory findings of the two studies mentioned above. In the study by Saffron and colleagues (2004), folate levels were similar between healthy and alcoholic patients, thus favoring the protective role of the polymorphism. In the study by Fabris and colleagues (2009), conversely, increased risk of HCC in people with the polymorphisms was only found in males, and not in females. Although the investigators did not measure folate levels, it is plausible that these levels would be lower in men, who tend to be heavier and more regular drinkers compared with women.

### Colorectal Cancer

Colorectal cancer is the third most common cancer and the second leading cause of cancer deaths for both sexes (Marin et al. 2012). Alcohol is a likely etiologic factor for this cancer (Baan et al. 2007). Several studies have indicated that epigenetic processes play a role in alcohol-related colorectal carcinogenesis. For example, rats chronically fed alcohol show genomic DNA hypomethylation but a normal pattern of methylation of the gene *TRP53*, which encodes a protein called p53, in the colonic mucosa (Choi et al. 1999). In a cohort of 609 patients, excessive alcohol use was associated with increased risk of colon cancer with global DNA hypomethylation (Schernhammer et al. 2010). In addition, people with low folate intake/high alcohol intake show a higher frequency of promoter methylation of genes involved in colorectal cancer carcinogenesis (e.g., *APC-1A*, *p14ARF*, *p16INK4a*, *hMLH1*, *O6-MGMT*, and *RASSF1A*) compared with people with high folate intake/low alcohol intake (Van Engeland et al. 2003). Many of these genes have fundamental roles in many cellular pathways, including DNA repair (e.g., *hMLH1*, *O6-MGMT*) and cell cycle

control (e.g., *p16INK4a*) (Portela and Esteller 2010).

The association between *MTHFR* polymorphisms and colon cancer has been studied extensively, and several factors have been found to influence the relation between the *MTHFR* variants and cancer risk (for a review, see Sharp and Little 2004). For example, one study (Iacopetta et al. 2009) has shown that colon cancer risk in homozygous carriers was dependent on the location of the cancer in the colon. The investigators observed that the TT genotype was associated with an increased risk for cancer in colon regions closer to the small intestine (i.e., proximal colon cancer) (adjusted odds ratio (AOR) = 1.29) but with a decreased risk for cancer in colon regions closer to the rectum (i.e., distal cancers) (AOR = 0.87). More importantly, the increased risk for proximal cancers was especially pronounced in individuals with high alcohol consumption (AOR = 1.90). Microsatellite instability also has been related with colon cancer risk in people with the TT genotype (Shannon et al. 2002). Finally, the dietary intake of folate and alcohol also has been associated with colon cancer risk in people carrying the *MTHFR* C variant. Thus, Kim and colleagues (2012) observed that low folate intake together with high alcohol intake increased the risk of colon cancer in people with either the CC or the CT genotype.

A subset of colorectal cancers exhibit promoter methylation in multiple genes; these tumors are referred to as the CpG island methylator phenotype (CIMP) (Toyota et al. 1999). The frequency of these tumors depends on the location of the cancer. Thus, 30 to 40 percent of sporadic proximal colon cancers are CIMP<sup>+</sup>, compared with 3 to 12 percent of distal colon and rectal cancers (Curtin et al. 2011). The presence of the A1298C *MTHFR* polymorphism, interacting with diet, may be involved in the development of highly CpG-methylated colon cancers. Homozygous and heterozygous genotypes in conjunction with a high-risk

dietary pattern (i.e., low folate and methionine intake and high alcohol use) were associated with CIMP<sup>+</sup> phenotype (Curtin et al. 2007).

### Breast Cancer

Breast cancer is the second leading cause of cancer death among women (DeSantis et al. 2011). Low doses of alcohol consumption (i.e.,  $\leq 1$  drink/day) increase the risk of breast cancer by about 4 percent (Hamajima et al. 2002), whereas heavy alcohol consumption (i.e.,  $\geq 3$  drink/day) is associated with an increase in risk of 40 to 50 percent (Pelucchi et al. 2011; Seitz

**Tobacco and alcohol are the major risk factors of upper aerodigestive tract cancers, or head and neck cancers, including cancers of the oral cavity, pharynx, larynx, and esophagus.**

et al. 2012). In addition, high frequency of alcohol consumption is associated with increased breast cancer mortality (Allemani et al. 2011). The role of epigenetic mechanisms in alcohol-related breast cancer also has been investigated. In a recent study of the methylation profiles of 1,413 CpG sites, Christensen and colleagues (2010) showed a strong trend toward decreased DNA methylation with increasing alcohol intake, and a trend toward increased methylation with increasing dietary folate. Other studies have shown altered methylation patterns for several genes associated with alcohol consumption, including hypermethylation of the *ER- $\alpha$*  (Zhu et al. 2003) and *E-cadherin* genes (Tao et al. 2011) and hypomethylation of *p16* (Tao et al. 2011).

As with other cancers, the person's genotype for the *C677T MTHFR* variant modulates the effect of alcohol

consumption on breast cancer risk. Thus, women with the TT genotype are at a higher risk of breast cancer than those with other genotypes. In postmenopausal women, the breast cancer risk was increased in women with the *C677T MTHFR* variant who had high lifetime daily alcohol intake, suggesting that folate metabolism has an impact on cancer development (Platek et al. 2009). As mentioned earlier, chronic alcohol abuse can cause folate deficiency, which is a well-documented risk factor for breast cancer (Sellers et al. 2001). Why this risk is only observed in post-menopausal women is not clear, but the levels of estrogen in the woman's body may play a role. Alcohol can interfere with estrogen pathways and increase the levels of estrogen in the blood (Dumitrescu and Shields 2005). Higher estrogen exposure, in turn, can induce aberrant DNA methylation associated with breast carcinogenesis both in vivo and in vitro (Fernandez and Russo 2010). These observations suggest another possible mechanism of alcohol-induced carcinogenesis, at least in breast cancer.<sup>2</sup>

### Upper Aerodigestive Tract Cancer

Tobacco and alcohol are the major risk factors of upper aerodigestive tract cancers, or head and neck cancers, including cancers of the oral cavity, pharynx, larynx, and esophagus. Each year, about 125,000 new cases of upper aerodigestive tract cancers are diagnosed across Europe, and more than half of the patients die from the disease (Anantharaman et al. 2011).

Several studies have shown an association between alcohol and aberrant DNA methylation in head and neck cancers and that the degree of DNA

<sup>2</sup> A woman's risk of breast cancer depends at least in part on her overall exposure to estrogen across her life span. Thus, women who begin menstruating earlier and reach menopause later have a higher cumulative estrogen exposure and, consequently, breast cancer risk than women with a shorter "menstrual life." Similarly, postmenopausal alcoholic women have higher estrogen levels than women who are not alcoholic, increasing their overall estrogen exposure, which may in turn increase breast cancer risk.



hypomethylation is associated with alcohol use in these cancers. A study examining the DNA methylation profiles of 1,413 CpG loci from 773 genes in head and neck squamous cell carcinomas showed that significant associations existed between methylation profiles and alcohol consumption (Marsit et al. 2009). Other small-scale studies also have shown alcohol-associated promoter hypermethylation for several genes, including *E-cadherin* and *p16INKa* (Hasegawa et al. 2002), p15 (Chang et al. 2004), *MGMT* (Puri et al. 2005), *p14ARF* (Ishida et al. 2005), *SFRP1* (Marsit et al. 2006), and *Fusel 18* and *Septin 9* (Bennett et al. 2010). The genotype at the *C677T MTHFR* variant also plays a role in the risk of alcohol-related upper aerodigestive tract cancers. Thus, heavy-drinking individuals with the TT genotype have an increased risk of oral cancer compared with the CC genotype (Supic et al. 2011) but a decreased risk of esophageal cancer (Yang et al. 2005). Thus, the effect of the *MTHFR* polymorphism appears to differ substantially depending on the type of cancer.

## Conclusions and Outlook

Aberrant DNA methylation is a hallmark of cancer development, and many studies have shown its contribution to tumor initiation and progression. In fact, methylation patterns nowadays are used as markers for cancer detection, tumor prognosis, and prediction of treatment responses. At the moment, evidence suggests that alcohol use is associated with aberrant DNA methylation patterns in several types of cancer. However, most studies have relied on looking at individual genes or a limited number of CpG loci. Genome-wide DNA methylation analyses may yield comprehensive maps of DNA methylation changes in alcohol-associated carcinogenesis, which could be important for use in pharmacoeugenetics, serving as additional markers for cancer detection, prognosis, and treatment response. Furthermore, despite all the

progress that has been made in elucidating how alcohol consumption might lead to altered DNA methylation patterns, the molecular mechanisms that lead to these alterations have to be better characterized so that effective therapies could be devised. ■

## Acknowledgements

This work is supported by NIH grants RO1-AT-1576, RO1-AT-004896, and RO1-DK-51719 (to S.C.L., M.L.M.-C., and J.M.M.); SAF 2011-29851 (to J.M.M.); ETORTEK-2011, Sanidad Gobierno Vasco 2008, Educación Gobierno Vasco 2011 (PI2011/29), and FIS (PI11/01588) (to M.L.M.-C.); FIS (PS09/00094), Fundación Científica de la Asociación Española Contra el Cáncer (Cancer Infantil), and the Program Ramón y Cajal, Spanish Ministry of Science and Innovation (to A.W.); and Sanidad Gobierno Vasco 2011 (to M.V.R.). CIBERehd is funded by the Instituto de Salud Carlos III.

## Financial Disclosure

The authors declare that they have no competing financial interests.

## References

- ALLEMANI, C.; BERRINO, F.; KROGH, V.; ET AL. Do pre-diagnostic drinking habits influence breast cancer survival? *Tumori* 97(2):142-148, 2011. PMID: 21617706
- ANANTHARAMAN, D.; MARRON, M.; LAGIU, P.; ET AL. Population attributable risk of tobacco and alcohol for upper aerodigestive tract cancer. *Oral Oncology* 47(8):725-731, 2011. PMID: 21684805
- BAAN, R.; STRAIF, K.; GROSSE, Y.; ET AL. Carcinogenicity of alcoholic beverages. *Lancet Oncology* 8(4):292-293, 2007. PMID: 17431955
- BARAK, A.J.; BECKENHAUER, H.C.; AND TUMA D.J. Betaine effects on hepatic methionine metabolism elicited by short-term ethanol feeding. *Alcohol* 13(5):483-486, 1996. PMID: 8888945
- BENNETT, K.L.; LEE, W.; LAMARRE, E.; ET AL. HPV status-independent association of alcohol and tobacco exposure or prior radiation therapy with promoter methylation of *FUSSEL18*, *EBF3*, *IRX1*, and *SEPT9*, but not *SLC5A8*, in head and neck squamous cell carcinomas. *Genes*,

*Chromosomes & Cancer* 49(4):319-326, 2010. PMID: 20029986

BIELAWSKI, D.M.; ZAHER, F.M.; SVINARICH, D.M.; AND ABEL, E.L. Paternal alcohol exposure affects sperm cytosine methyltransferase messenger RNA levels. *Alcoholism: Clinical and Experimental Research* 26(3):347-351, 2002. PMID: 11923587

BOFFETTA, P.; HASHIBE, M.; LA VECCHIA, C.; ET AL. The burden of cancer attributable to alcohol drinking. *International Journal of Cancer* 119(4):884-887, 2006. PMID: 16557583

BÖNSCH, D.; LENZ, B.; FISZER, R.; ET AL. Lowered DNA methyltransferase (DNMT-3b) mRNA expression is associated with genomic DNA hypermethylation in patients with chronic alcoholism. *Journal of Neural Transmission* 113(9):1299-1304, 2006. PMID: 16463117

CALVISI, D.F.; LADU, S.; GORDEN, A.; ET AL. Mechanistic and prognostic significance of aberrant methylation in the molecular pathogenesis of human hepatocellular carcinoma. *Journal of Clinical Investigation* 117(9):2713-2722. PMID: 17717605

CHANG, H.W.; LING, G.S.; WEI, W.I.; AND YUEN, A.P. Smoking and drinking can induce p15 methylation in the upper aerodigestive tract of healthy individuals and patients with head and neck squamous cell carcinoma. *Cancer* 101(1):125-132, 2004. PMID: 15221997

CHOI, S.W., AND FRISO, S. Interactions between folate and aging for carcinogenesis. *Clinical Chemistry and Laboratory Medicine* 43(10):1151-1157, 2005. PMID: 16197313

CHOI, S.W.; STICKEL, F.; BAIK, H.W.; ET AL. Chronic alcohol consumption induces genomic but not p53-specific DNA hypomethylation in rat colon. *Journal of Nutrition* 129(11):1945-50, 1999. PMID: 10539767

CHRISTENSEN, B.C.; KELSEY, K.T.; ZHENG, S.; ET AL. Breast cancer DNA methylation profiles are associated with tumor size and alcohol and folate intake. *PLoS Genetics* 6(7):e1001043, 2010. PMID: 20686660

CRAVO, M.L.; GLÓRIA, L.M.; SELHUB, J.; ET AL. Hyperhomocysteinemia in chronic alcoholism: Correlation with folate, vitamin B-12, and vitamin B-6 status. *American Journal of Clinical Nutrition* 63(2):220-224, 1996. PMID: 8561063

CURTIN, K.; SLATTERY, M.L.; AND SAMOWITZ, W.S. CpG island methylation in colorectal cancer: Past, present and future. *Pathology Research International* 2011:902674, 2011. PMID: 21559209

CURTIN, K.; SLATTERY, M.L.; ULRICH, C.M.; ET AL. Genetic polymorphisms in one-carbon metabolism: Associations with CpG island methylator phenotype (CIMP) in colon cancer and the modifying effects of diet. *Carcinogenesis* 28(8):1672-1679, 2007. PMID: 17449906

DESANTIS, C.; SIEGEL, R.; BANDI, P.; AND JEMAL, A. Breast cancer statistics, 2011. CA: *A Cancer Journal for Clinicians* 61(6):409-418, 2011. PMID: 21969133

DUMITRESCU, R.G., AND SHIELDS, P.G. The etiology of alcohol-induced breast cancer. *Alcohol* 35(3):213-25, 2005. PMID: 16054983.

EDAMOTO, Y.; HARA, A.; BIERNAT, W.; ET AL. Alterations of RB1, p53 and Wnt pathways in hepatocellular carcinoma.

mas associated with hepatitis C, hepatitis B and alcoholic liver cirrhosis. *International Journal of Cancer* 106(3):334–341, 2003. PMID: 12845670

ESTELLER, M. Epigenetics in cancer. *New England Journal of Medicine* 358(11):1148–1159, 2008. PMID: 18337604

FABRIS, C.; TONIUTTO, P.; FALLETTI, E.; ET AL. MTHFR C677T polymorphism and risk of HCC in patients with liver cirrhosis: Role of male gender and alcohol consumption. *Alcoholism: Clinical and Experimental Research* 33(1):102–107, 2009. PMID: 18945219

FERNANDEZ, S.V., AND RUSSO, J. Estrogen and xenoestrogens in breast cancer. *Toxicologic Pathology* 38(1):110–122, 2010. PMID: 19933552

GARRO, A.J.; McBETH, D.L.; LIMA, V.; AND LIEBER, C.S. Ethanol consumption inhibits fetal DNA methylation in mice: Implications for the fetal alcohol syndrome. *Alcoholism: Clinical and Experimental Research* 15(3):395–398, 1991. PMID: 1877725

GIOVANNUCCI, E.; RIMM, E.B.; ASCHERIO, A.; ET AL. Alcohol, low-methionine—low-folate diets, and risk of colon cancer in men. *Journal of the National Cancer Institute* 87(4):265–273, 1995. PMID: 7707417

HALSTED, C.H.; ROBLES, E.A.; AND MEZEY, E. Decreased jejunal uptake of labeled folic acid (3H-PGA) in alcoholic patients: Roles of alcohol and nutrition. *New England Journal of Medicine* 285(13):701–706, 1971. PMID: 5571128

HALSTED, C.H.; VILLANUEVA, J.; CHANDLER, C.J.; ET AL. Ethanol feeding of micropigs alters methionine metabolism and increases hepatocellular apoptosis and proliferation. *Hepatology* 23(3):497–505, 1996. PMID: 8617429

HAMAJIMA, N.; HIROSE, K.; TAJIMA, K.; ET AL. Alcohol, tobacco and breast cancer—Collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *British Journal of Cancer* 87(11):1234–1245, 2002. PMID: 12439712

HAMID, A. Folate malabsorption and its influence on DNA methylation during cancer development. *DNA and Cell Biology* April 2, 2012. [Epub ahead of print]. PMID: 22468673

HASEGAWA, M.; NELSON, H.H.; PETERS, E.; ET AL. Patterns of gene promoter methylation in squamous cell cancer of the head and neck. *Oncogene* 21(27):4231–4236, 2002. PMID: 12082610

HERNANDEZ-VARGAS, H.; LAMBERT, M.P.; LE CALVEZ-KELM, F.; ET AL. Hepatocellular carcinoma displays distinct DNA methylation signatures with potential as clinical predictors. *PLoS One* 5(3):e9749, 2010. PMID: 20305825

IACOPETTA, B.; HEYWORTH, J.; GIRSCHIK, J.; ET AL. The MTHFR C677T and DeltaDNMT3B C-149T polymorphisms confer different risks for right- and left-sided colorectal cancer. *International Journal of Cancer* 125(1):84–90, 2009. PMID: 19326430

ISHIDA, E.; NAKAMURA, M.; IKUTA, M.; ET AL. Promotor hypermethylation of p14ARF is a key alteration for progression of oral squamous cell carcinoma. *Oral Oncology* 41(6):614–622, 2005. PMID: 15975525

KIM, J.; CHO, Y.A.; KIM, D.H.; ET AL. Dietary intake of folate and alcohol, MTHFR C677T polymorphism, and colorec-

tal cancer risk in Korea. *American Journal Clinical Nutrition* 95(2):405–412, 2012. PMID: 22218157

KIM, Y.I. Nutritional epigenetics: Impact of folate deficiency on DNA methylation and colon cancer susceptibility. *Journal of Nutrition* 135(11):2703–2709, 2005. PMID: 16251634

KOBOR, M.S., AND WEINBERG, J. Epigenetics and fetal alcohol spectrum disorders. *Alcohol Research & Health* 34(1):29–37.

LAMBERT, M.P.; PALIWAL, A.; VAISSIÈRE, T.; ET AL. Aberrant DNA methylation distinguishes hepatocellular carcinoma associated with HBV and HCV infection and alcohol intake. *Journal of Hepatology* 54(4):705–715, 2011. PMID: 21146512

LARSSON, S.C.; GIOVANNUCCI, E.; AND WOLK, A. Vitamin B6 intake, alcohol consumption, and colorectal cancer: A longitudinal population-based cohort of women. *Gastroenterology* 128(7):1830–1837, 2005. PMID: 15940618

LEE, T.D.; SADDA, M.E.; MENDLER, M.H.; ET AL. Abnormal hepatic methionine and glutathione metabolism in patients with alcoholic hepatitis. *Alcoholism: Clinical and Experimental Research* 28(1):173–181, 2004. PMID: 14745316

LU, S.C. S-Adenosylmethionine. *International Journal of Biochemistry & Cell Biology* 32(4):391–395, 2000. PMID: 10762064

LU, S.C., AND MATO, J.M. S-Adenosylmethionine in cell growth, apoptosis and liver cancer. *Journal of Gastroenterology and Hepatology* 23(Suppl. 1):S73–S77, 2008. PMID: 18336669

MA, J.; STAMPFER, M.J.; GIOVANNUCCI, E.; ET AL. Methylenetetrahydrofolate reductase polymorphism, dietary interactions, and risk of colorectal cancer. *Cancer Research* 57(6):1098–1102, 1997. PMID: 9067278

MARIN, J.J.; SANCHEZ DE MEDINA, F.; CASTAÑO, B.; ET AL. Chemoprevention, chemotherapy, and chemoresistance in colorectal cancer. *Drug Metabolism Reviews* 44(2):148–172, 2012. PMID: 22497631

MARSIT, C.J.; CHRISTENSEN, B.C.; HOUSEMAN, E.A.; ET AL. Epigenetic profiling reveals etiologically distinct patterns of DNA methylation in head and neck squamous cell carcinoma. *Carcinogenesis* 30(3):416–422, 2009. PMID: 19126652

MARSIT, C.J.; McCLEAN, M.D.; FURNISS, C.S.; AND KELSEY, K.T. Epigenetic inactivation of the SFRP genes is associated with drinking, smoking and HPV in head and neck squamous cell carcinoma. *International Journal of Cancer* 119(8):1761–1766, 2006. PMID: 16708382

MATO, J.M.; MARTÍNEZ-CHANTAR, M.L.; AND LU, S.C. Methionine metabolism and liver disease. *Annual Review of Nutrition* 28:273–293, 2008. PMID: 18331185

MOGHE, A.; JOSHI-BARVE, S.; GHARE, S.; ET AL. Histone modifications and alcohol-induced liver disease: Are altered nutrients the missing link? *World Journal of Gastroenterology* 17(20):2465–2472, 2011. PMID: 21633651

MOTULSKY, A.G. Nutritional ecogenetics: Homocysteine-related arteriosclerotic vascular disease, neural tube

defects, and folic acid. *American Journal Human Genetics* 58(1):17–20, 1996. PMID: 8554053

PELUCCI, C.; TRAMACERE, I.; BOFFETTA, P.; ET AL. Alcohol consumption and cancer risk. *Nutrition and Cancer* 63(7):983–990, 2011. PMID: 21864055

PLATEK, M.E.; SHIELDS, P.G.; MARIAN, C.; ET AL. Alcohol consumption and genetic variation in methylenetetrahydrofolate reductase and 5-methyltetrahydrofolate-homocysteine methyltransferase in relation to breast cancer risk. *Cancer Epidemiology, Biomarkers & Prevention* 18(9):2453–2459, 2009. PMID: 19706843

PORTELA, A., AND ESTELLER, M. Epigenetic modifications and human disease. *Nature Biotechnology* 28(10):1057–1068, 2010. PMID: 20944598

PURI, S.K.; SI, L.; FAN, C.Y.; AND HANNA, E. Aberrant promoter hypermethylation of multiple genes in head and neck squamous cell carcinoma. *American Journal of Otolaryngology* 26(1):12–17, 2005. PMID: 15635575

REHM, J.; ROOM, R.; MONTEIRO, M.; ET AL. Alcohol use. In: Ezzati, M.; Murray, C.; Lopez, A.D.; and Rodgers, A., Eds. *Comparative Quantification of Health Risks: Global and Regional Burden of Disease Attributable to Selected Major Risk Factors*. Geneva, Switzerland: World Health Organization, 2004, pp. 959–1108.

RODRÍGUEZ-PAREDES, M., AND ESTELLER M. Cancer epigenetics reaches mainstream oncology. *Nature Medicine* 17(3):330–339, 2011. PMID: 21386836

ROSS, S.A., AND POIRIER, L. Proceedings of the Trans-HHS Workshop: Diet, DNA methylation processes and health. *Journal of Nutrition* 132(Suppl. 8):2329S–2332S, 2002. PMID: 12163686

SAFFROY, R.; PHAM, P.; CHIAPPINI, F.; ET AL. The MTHFR 677C > T polymorphism is associated with an increased risk of hepatocellular carcinoma in patients with alcoholic cirrhosis. *Carcinogenesis* 25(8):1443–1448, 2004. PMID: 15033905

SCHERNHAMMER, E.S.; GIOVANNUCCI, E.; KAWASAKI, T.; ET AL. Dietary folate, alcohol and B vitamins in relation to LINE-1 hypomethylation in colon cancer. *Gut* 59(6):794–799, 2010. PMID: 19828464

SEITZ, H.K.; PELUCCI, C.; BAGNARDI, V.; AND LA VECCHIA, C. Epidemiology and pathophysiology of alcohol and breast cancer: Update 2012. *Alcohol and Alcoholism* 47(3):204–212, 2012. PMID: 22459019

SEITZ, H.K., AND STICKEL, F. Molecular mechanisms of alcohol-mediated carcinogenesis. *Nature Reviews. Cancer* 7(8):599–612, 2007. PMID: 17646865

SELLERS, T.A.; KUSHI, L.H.; CERHAN, J.R.; ET AL. Dietary folate intake, alcohol, and risk of breast cancer in a prospective study of postmenopausal women. *Epidemiology* 12(4):420–428, 2001. PMID: 11416780

SHANNON, B.; GNANASAMPANTHAN, S.; BEILBY, J.; AND IACOPETTA, B. A polymorphism in the methylenetetrahydrofolate reductase gene predisposes to colorectal cancers with microsatellite instability. *Gut* 50(4):520–524, 2002. PMID: 11889073

SHARP, L., AND LITTLE, J. Polymorphisms in genes involved in folate metabolism and colorectal neoplasia: A HuGE review. *American Journal of Epidemiology* 159(5):423–443, 2004. PMID: 14977639

- SUPIC, G.; JOVIC, N.; KOZOMARA, R.; ET AL. Interaction between the MTHFR C677T polymorphism and alcohol—Impact on oral cancer risk and multiple DNA methylation of tumor-related genes. *Journal of Dental Research* 90(1):65–70, 2011. PMID: 20940365
- TAO, M.H.; MARIAN, C.; SHIELDS, P.G.; ET AL. Alcohol consumption in relation to aberrant DNA methylation in breast tumors. *Alcohol* 45(7):689–699, 2011. PMID: 21168302
- TISCHOFF, I., AND TANNAPPE, A. DNA methylation in hepatocellular carcinoma. *World Journal of Gastroenterology* 14(11):1741–1748, 2008. PMID: 18350605
- TOYOTA, M., AND ISSA J.P. CpG island methylator phenotypes in aging and cancer. *Seminars in Cancer Biology* 9(5):349–357, 1999. PMID: 10547343
- TSUKAMOTO, H., AND LU, S.C. Current concepts in the pathogenesis of alcoholic liver injury. *FASEB Journal* 15(8):1335–1349, 2001. PMID: 11387231
- VAN DER PUT, N.M.; STEEGERS-THEUNISSEN, R.P.; FROSST, P.; ET AL. Mutated methylenetetrahydrofolate reductase as a risk factor for spina bifida. *Lancet* 346(8982):1070–1071, 1995. PMID: 7564788
- VAN DER PUT, N.M.; VAN STRAATEN, H.W.; TRIJBELS, F.J.; AND BLOM, H.J. Folate, homocysteine and neural tube defects: An overview. *Experimental Biology and Medicine (Maywood)* 226(4):243–270, 2001. PMID: 11368417
- VAN ENGELAND, M.; WEIJENBERG, M.P.; ROEMEN, G.M.; ET AL. Effects of dietary folate and alcohol intake on promoter methylation in sporadic colorectal cancer: The Netherlands cohort study on diet and cancer. *Cancer Research* 63(12):3133–3137, 2003. PMID: 12810640
- YANG, C.X.; MATSUO, K.; ITO, H.; ET AL. Gene-environment interactions between alcohol drinking and the MTHFR C677T polymorphism impact on esophageal cancer risk: Results of a case-control study in Japan. *Carcinogenesis* 26(7):1285–1290, 2005. PMID: 15790587
- YUAN, J.M.; LU, S.C.; VAN DEN BERG, D.; ET AL. Genetic polymorphisms in the methylenetetrahydrofolate reductase and thymidylate synthase genes and risk of hepatocellular carcinoma. *Hepatology* 46(3):749–758, 2007. PMID: 17659576
- ZHU, K.; DAVIDSON, N.E.; HUNTER, S.; ET AL. Methyl-group dietary intake and risk of breast cancer among African-American women: A case-control study by methylation status of the estrogen receptor alpha genes. *Cancer Causes & Control* 14(9):827–836, 2003. PMID: 14682440

# Announcing Two New Alcohol Alerts



Number 84

## The Genetics of Alcoholism

Why can some people have a glass of wine or beer with their meal without feeling compelled to drink more, whereas others can't seem to stop drinking? Can some people “hold” their liquor better than others? Does alcoholism tend to run in families? Scientists have been trying to find answers to questions such as these for several decades, seeking to identify the factors that influence a person's risk of becoming alcohol dependent.

This *Alcohol Alert* explores the genetics of alcoholism and provides an update on what we know about how hereditary influences drinking.



Number 85

## The Link Between Stress and Alcohol

A new job, a death in the family, moving across the country, a breakup, or getting married—all are situations that can result in psychological and physical symptoms collectively known as “stress.” One way that people may choose to cope with stress is by turning to alcohol.

This *Alcohol Alert* explores the relationship between alcohol and stress, including identifying some common sources of stress, examining how the body responds to stressful situations, and the role that alcohol plays—both in alleviating and perpetuating stress.

<http://pubs.niaaa.nih.gov/publications/>