

Structured Clinical Interview for the DSM (SCID) Substance Use Disorder Module—Adapted for Adolescents

BRIEF DESCRIPTION	This is a DSM-IV substance use disorder interview adapted for use with adolescents.
TARGET POPULATION	<input type="checkbox"/> Adults <input checked="" type="checkbox"/> Adolescents (over 16 years) Groups for which this instrument might be especially helpful? Adolescents suspected of having an alcohol use disorder or another substance use disorder
ADMINISTRATIVE ISSUES	Number of items: <i>Varies</i> Number of subscales: Format(s): <input type="checkbox"/> Pencil-and-paper self-administered <input checked="" type="checkbox"/> Interview (structured) <input type="checkbox"/> Observation <input type="checkbox"/> Computer self-administered <input type="checkbox"/> Other Time required for administration: <i>30 to 90 minutes</i> Administered by: <i>Trained clinician</i> Training required for administration? <input checked="" type="checkbox"/> yes <input type="checkbox"/> no
SCORING	Time required to score/interpret: <i>10 to 15 minutes</i> Scored by: <i>Interviewer</i> Scoring key? <input checked="" type="checkbox"/> yes <input type="checkbox"/> no Computerized scoring or interpretation available? <input type="checkbox"/> yes <input checked="" type="checkbox"/> no Norms available? <input type="checkbox"/> yes <input checked="" type="checkbox"/> no Instrument normed on subgroups? <input type="checkbox"/> yes <input checked="" type="checkbox"/> no

PSYCHOMETRICS

Have reliability studies been done? yes no

What measure(s) of reliability was used?

Test-retest

Split half

Internal consistency

Have validity studies been done? yes no

What measures of validity have been derived?

Content

Criterion (predictive, concurrent, "postdictive")

Construct

CLINICAL UTILITY OF INSTRUMENT

Case identification and treatment planning

RESEARCH APPLICABILITY

Composition of research groups and assessment of outcomes

SOURCE, COST AND COPYRIGHT ISSUES

Copyright: yes no

Cost: *Public domain*

Source: Chris Martin, Ph.D.
Western Psychiatric Institute
Pittsburgh, PA

SOURCE REFERENCE

Martin, C., Kaczynski, N., Maisto, S., Bukstein, O. & Moss, H. (1995). Patterns of DSM-IV alcohol abuse and dependence symptoms in adolescent drinkers. *Journal of Studies on Alcohol*, 56, 672-680.

SUPPORTING REFERENCES

Martin, C., Pollock, N., Bukstein, O. & Lynch, K. (2000). Inter-rater reliability of the SCID alcohol and substance use disorders section among adolescents. *Drug and Alcohol Dependence*, 59, 173-176. Erratum: (2000) *Drug and Alcohol Dependence*, 60, 323.

Martin, C., Langenbucher, J., Kaczynski, N. & Chung, T. (1996). Staging in the onset of DSM-IV alcohol abuse and dependence symptoms in adolescents: Survival/hazard analyses. *Journal of Studies on Alcohol*, 57, 549-558.

Martin, C., Kaczynski, N., Maisto, S. & Tarter, R. (1996). Polydrug use in adolescent drinkers with and without DSM-IV alcohol abuse and dependence. *Alcoholism: Clinical and Experimental Research*, 20, 1099-1108.